

Near threshold $p\bar{p}$ enhancement in B and J/Ψ decay

J. Haidenbauer¹, Ulf-G. Meißner^{1,2}, A. Sibirtsev²

¹*Institut für Kernphysik (Theorie), Forschungszentrum Jülich, D-52425 Jülich, Germany*

²*Helmholtz-Institut für Strahlen- und Kernphysik (Theorie),
Universität Bonn, Nußallee 14-16, D-53115 Bonn, Germany*

The near-threshold enhancement in the $p\bar{p}$ invariant mass spectrum from the $B^+ \rightarrow K^+ p\bar{p}$ decay reported recently by the BaBar Collaboration is studied within the Jülich $N\bar{N}$ model. We illustrate that the invariant mass dependence of the $p\bar{p}$ spectrum close to the threshold can be reproduced by the final state interactions. This explanation is in line with our previous analysis of the $p\bar{p}$ invariant mass spectrum from the $J/\Psi \rightarrow \gamma p\bar{p}$ decay measured by the BES Collaboration. We also comment on a structure found recently in the $\pi^+ \pi^- \eta'$ mass spectrum of the radiative J/Ψ decay by the BES Collaboration. In particular we argue that one should be rather cautious in bringing this structure in connection with the enhancement found in the $p\bar{p}$ invariant mass spectrum or with the existence of $N\bar{N}$ bound states.

PACS numbers: 11.80.-m; 13.60.Le; 13.75.Jz; 14.65.Dw; 25.80.Nv

A first indication for a near-threshold enhancement in the proton-antiproton $p\bar{p}$ invariant mass spectrum from the $B^+ \rightarrow K^+ p\bar{p}$ and $\bar{B}^0 \rightarrow D^0 p\bar{p}$ decays were reported by the Belle Collaboration [1, 2]. Soon afterwards a much more significant evidence of a $p\bar{p}$ enhancement, i.e. with high statistics and high mass resolution, was observed by the BES Collaboration [3] in the reaction $J/\Psi \rightarrow \gamma p\bar{p}$. More recently the Belle Collaboration [4] found also a near-threshold $p\bar{p}$ enhancement in the decays $B^+ \rightarrow \pi^+ p\bar{p}$, $B^+ \rightarrow K^+ p\bar{p}$, $B^0 \rightarrow K^0 p\bar{p}$ and $B^+ \rightarrow K^{*+} p\bar{p}$, while the CLEO Collaboration detected such an enhancement in (the unsubtracted) data for $\Upsilon(1S) \rightarrow \gamma p\bar{p}$ [5]. However, in all these cases the results are marred by low statistics. Very recently the BaBar Collaboration presented a new and high statistics measurement of the $B^+ \rightarrow K^+ p\bar{p}$ decay [6] confirming the threshold peaking in the $p\bar{p}$ invariant mass, see also [7].

The high statistics data by the BES Collaboration triggered a number of theoretical speculations where the observed enhancement in the invariant $p\bar{p}$ mass spectrum was interpreted as evidence for a $p\bar{p}$ bound state or baryonium [8, 9], or for exotic glueball states [10, 11]. Alternatively, we [12] but also others [13, 14, 15, 16] demonstrated that the near-threshold enhancement in the $p\bar{p}$ invariant mass spectrum from the $J/\Psi \rightarrow \gamma p\bar{p}$ decay could be simply due to the final state interactions (FSI) between the outgoing proton and antiproton. Specifically, our calculation based on the realistic Jülich $N\bar{N}$ model [17, 18] and the one by Loiseau and Wycech [16], utilizing the Paris $N\bar{N}$ model, explicitly confirmed the significance of FSI effects estimated in the initial studies [13, 14, 15] within the effective range approximation.

In the present paper we want to investigate whether the near-threshold enhancement in the $p\bar{p}$ invariant mass spectrum, visible in the high statistics data on the reaction $B^+ \rightarrow K^+ p\bar{p}$, can likewise be understood in terms of the $p\bar{p}$ FSI. In our study of the $J/\Psi \rightarrow \gamma p\bar{p}$ decay we considered the $p\bar{p}$ FSI interaction in the 1S_0 and 3P_0 partial waves and the $I = 0$ and $I = 1$ isospin channels. Other $p\bar{p}$ S - and P -waves are ruled out by conservation

laws for parity, charge-conjugation and total angular momentum together with the measured photon angular distribution from the $J/\Psi \rightarrow \gamma p\bar{p}$ decay, which agrees with that expected from the $p\bar{p}$ state being in both 1S_0 and 3P_0 states. We found that the mass dependence of the $p\bar{p}$ spectrum close to the threshold can be reproduced by the S -wave $p\bar{p}$ FSI in the isospin $I = 1$ state. In case of the $B^+ \rightarrow K^+ p\bar{p}$, the weak interaction is involved. As a consequence, the selection rules are less rigid and now other $p\bar{p}$ S - and P -waves are allowed too and could produce FSI effects in the near-threshold region. Thus, besides the effects resulting from the 1S_0 and 3P_0 partial waves we explore here also those of the 3S_1 and 3P_1 states.

Like in our earlier paper we utilize the total spin-averaged (dimensionless) $B^+ \rightarrow K^+ p\bar{p}$ reaction amplitude A and not directly the measured $p\bar{p}$ invariant mass spectrum, because that allows us to get rid of trivial kinematical factors. The $B^+ \rightarrow K^+ p\bar{p}$ decay rate is given in terms of A by [19]

$$d\Gamma = \frac{|A|^2}{2^9 \pi^5 m_{B^+}^2} \lambda^{1/2}(m_{B^+}^2, M^2, m_{K^+}) \times \lambda^{1/2}(M^2, m_p^2, m_{\bar{p}}^2) dM d\Omega_p d\Omega_K, \quad (1)$$

where the Kallen function λ is defined by $\lambda(x, y, z) = ((x - y - z)^2 - 4yz)/4x$, $M \equiv M(p\bar{p})$ is the invariant mass of the $p\bar{p}$ system, Ω_p is the proton angle in that system, while Ω_K is the K^+ angle in the B^+ rest frame. After averaging over the spin states and integrating over the angles, the differential decay rate is

$$\frac{d\Gamma}{dM} = \frac{\lambda^{1/2}(m_{B^+}^2, M^2, m_{K^+}) \sqrt{M^2 - 4m_p^2}}{2^6 \pi^3 m_{B^+}^2} |A|^2. \quad (2)$$

We use Eq. (2) for extracting $|A|^2$ from the data of the BaBar Collaboration. The corresponding results are shown in Fig. 1 by the filled circles.

We assume again the validity of the Watson-Migdal approach for the treatment of the FSI effect. It suggests




FIG. 1: Invariant $B^+ \rightarrow K^+ p\bar{p}$ amplitude $|A|^2$ as a function of the $p\bar{p}$ mass. The circles represent the experimental values of $|A|^2$ extracted from the BaBar data [6] via Eq. (2). The curves are the scattering amplitude squared ($|T|^2$) predicted by the $N\bar{N}$ model A(OBE) [17] for the 1S_0 , 3S_1 , 3P_0 , and 3P_1 partial waves and the $I=0$ (solid) and $I=1$ (dashed) channels, respectively. Note that the latter results have been normalized to $|A|^2$ at $M(p\bar{p}) - 2m_p = 50$ MeV. For comparison reasons we show also the corresponding results extracted from the data on $J/\Psi \rightarrow \gamma p\bar{p}$ [3] (open circles).

that the reaction amplitude for a production and/or decay reaction that is of short-ranged nature can be factorized in terms of an elementary (basically constant) production amplitude and the $p\bar{p}$ scattering amplitude T of the particles in the final state so that

$$A(M(p\bar{p})) \approx N \cdot T(M(p\bar{p})), \quad (3)$$

(cf. Ref. [12] for further details). Thus, we compare the extracted amplitude $|A|^2$ with the suitably normalized scattering amplitudes $|T|^2$ that result from the Jülich $N\bar{N}$ model. The curves shown in Fig. 1 correspond to the $p\bar{p}$ scattering amplitude squared calculated for the 1S_0 , 3S_1 , 3P_0 and 3P_1 partial waves, where the solid lines are the results for the isospin $I=0$ channel, while the dashed lines are for the $I=1$ channel.

As can be seen from Fig. 1, the enhancement of the near-threshold $p\bar{p}$ invariant mass spectrum from the $B^+ \rightarrow K^+ p\bar{p}$ decay, as observed in the new BaBar experiment [6], is fully in line with our previous results [12]. Fig. 1 contains also the $J/\Psi \rightarrow \gamma p\bar{p}$ reaction amplitude,

evaluated via

$$\frac{d\Gamma}{dM} = \frac{(m_{J/\Psi}^2 - M^2) \sqrt{M^2 - 4m_p^2}}{2^7 \pi^3 m_{J/\Psi}^3} |A_{J/\Psi}|^2, \quad (4)$$

again suitably normalized to facilitate an easy comparison of the dependence on the $p\bar{p}$ invariant mass. Obviously the data from both considered decay reactions are in reasonable agreement as far as the dependence on $M(p\bar{p})$ is concerned. As already mentioned, while for the $J/\Psi \rightarrow \gamma p\bar{p}$ data the final $p\bar{p}$ system is restricted to the 1S_0 and 3P_0 partial waves in the near-threshold region, due to selection rules and the measured photon angular distribution [3], the $B^+ \rightarrow K^+ p\bar{p}$ reaction allows also for other partial waves. But also here a measurement of the K^+ angular distribution could clarify whether the 1S_0 or 3P_0 partial waves are responsible for the $p\bar{p}$ enhancement, as for the J/Ψ , or rather the 3S_1 , 3P_1 or 1P_1 states. Conservation of the total angular momentum requires the K^+ to be either in a relative s wave to the $p\bar{p}$ system (for the 1S_0 or 3P_0 partial waves) or in a p wave (for 3S_1 , 3P_1 or 1P_1). We note that the invariant amplitude for the 1P_1 wave looks very similar to the one for 3P_1 .

Recently the CLEO Collaboration published results on the radiative decays of the $\Upsilon(1S)(9460)$ to the $p\bar{p}$ system [5]. Interestingly, also in this reaction one can see an enhancement in the $p\bar{p}$ invariant mass spectrum near threshold, cf. Fig. 6 in that paper. The authors presented also results of a reference measurement for the reaction $e^+e^- \rightarrow \gamma p\bar{p}$ at the energy $\sqrt{s} = 10.56$ GeV where a similar near-threshold enhancement in the $p\bar{p}$ mass spectrum is detected. We do not show corresponding results here because the accuracy and the mass resolution of those data is too low for allowing a meaningful comparison. However, we would like to comment on a conclusion drawn in Ref. [5]. In order to remove possible continuum background contributions the CLEO Collaboration subtracted the (scaled) $e^+e^- \rightarrow \gamma p\bar{p}$ mass spectrum from the one measured for the $\Upsilon(1S)$ radiative decay. The “corrected” $\Upsilon(1S) \rightarrow p\bar{p}$ data do not show an enhancement in the $p\bar{p}$ spectrum anymore. We believe that this is not surprising and, in fact, must be expected if the near-threshold enhancement comes indeed from the FSI in the $p\bar{p}$ system. Then, the same or a similar FSI must be present in $e^+e^- \rightarrow \gamma p\bar{p}$ as well as in $\Upsilon(1S) \rightarrow \gamma p\bar{p}$ and it must cancel to a large degree in a subtraction like the one performed in Ref. [5]. Accordingly, from our point of view there is no contradiction between the results of CLEO and those of the BES Collaboration for $J/\Psi \rightarrow \gamma p\bar{p}$ as suggested in Ref. [5]. Rather the CLEO results even strengthen the conjecture that the near-threshold enhancement in the $p\bar{p}$ spectrum seen in J/Ψ , B^+ , $\Upsilon(1S)$ etc. decays is due to the $p\bar{p}$ FSI.

Next, let us reflect on the present results in view of the earlier consideration concerning the J/Ψ decays. In our work on the $J/\Psi \rightarrow \gamma p\bar{p}$ spectrum we admitted that, because of our poor knowledge of the $N\bar{N}$ interaction near threshold and for some other reasons [12], explana-

tions for the enhancement other than final state interactions cannot be ruled out at the present stage. Specifically, we discussed [12] that intermediate pseudoscalar ($J^{PC}=0^{-+}$) meson resonances, for instance the $\pi(1800)$ resonance but also the $\eta(1760)$ [20], could couple to the $p\bar{p}$ channel and thus could play a role. In fact, we showed that the presence of these resonances in the decay $J/\Psi \rightarrow \gamma p\bar{p}$ is, in principle, in line with the BES data [3] once FSI effects are taken into account.

In this context it is interesting to note that recently the BES Collaboration reported [21] a resonance in the $J/\Psi \rightarrow \gamma \pi^+ \pi^- \eta'$ mode with mass 1833.7 ± 6.1 MeV and width of 67.7 ± 20.3 MeV obtained by fitting a Breit-Wigner function to the $\pi^+ \pi^- \eta'$ invariant spectrum. This resonance was denoted as a new $X(1835)$ state, arguing that it is not compatible with any of the meson resonance listed in Ref. [20].

Following our prescription [12] the authors of Ref. [21] re-fitted the $J/\Psi \rightarrow \gamma p\bar{p}$ spectrum including a pseudoscalar resonance and the $I = 0$ 1S_0 $p\bar{p}$ FSI of the Jülich $N\bar{N}$ model A(OBE) [17]. The fit yielded a mass of 1831 ± 7 MeV and a $\Gamma < 153$ MeV and led them to the conclusion that the $X(1835)$ properties as found in $J/\Psi \rightarrow \gamma \pi^+ \pi^- \eta'$ are consistent with expectations for the state that produces the strong $p\bar{p}$ mass threshold enhancement observed in the $J/\Psi \rightarrow \gamma p\bar{p}$ decay [21].

Though the authors of Ref. [21] admitted in the summary that other possible interpretations of the $X(1835)$ that have no relation to the observed near-threshold $p\bar{p}$

enhancement are not excluded, we think its worthwhile to elaborate further on that issue. Dividing the average total $J/\Psi \rightarrow \gamma p\bar{p}$ reaction amplitude $|A_{J/\Psi}|^2$, which is related to the differential decay rate via Eq. (4), by the $p\bar{p}$ scattering amplitude $|T|^2$ [12, 17] one can easily obtain the FSI corrected data as a function of the $p\bar{p}$ invariant mass $M(p\bar{p})$, cf. Fig. 2. Here we use the 1S_0 scattering amplitudes in the $I = 0$ and $I = 1$ isospin states.

In our experience the data do not allow to fix uniquely the resonance properties if the mass, width and the strength of the resonant contribution are unknown. In order to illustrate that we show the squared Breit-Wigner amplitudes for the $X(1835)$, the $\eta(1760)$ and $\pi(1800)$ with their properties given by the BES Collaboration [21] and the PDG [20], respectively. We only vary the coupling strength of the 0^{-+} state to the $p\bar{p}$ channel in order to reproduce the BES data [3]. Since the $X(1835)$ might be a $I^G(J^{PC})=0^+(0^{-+})$ resonance [21] one can certainly speculate whether it is the same object as the $\eta(1760)$ listed by the PDG [20]. In fact, the $\eta(1760)$ was also established in radiative J/Ψ decays, namely in the reaction $J/\Psi \rightarrow \gamma \rho \rho$ [22]. While the PDG cites only an averaged value, a glance into the original paper [22] makes clear that the mass as well as the width of the $\eta(1760)$ could not be reliably established from the data. Indeed, one of the six solutions with comparable χ^2 given in Ref. [22] (in Table IV) yields a mass and width of 1807 ± 10 MeV and 94 ± 12 MeV, which is not that far away from the values obtained by the BES Collaboration. We show also results based on the above resonance parameters in Fig. 2. It is obvious that there is practically no difference between the $X(1835)$ of Ref. [21] and the $\eta(1800)$ of Ref. [22] as far as the description of the $p\bar{p}$ invariant mass spectrum is concerned.

Anyway, the properties of the $X(1835)$ are indeed consistent with the measured near-threshold enhancement in the $p\bar{p}$ spectrum of the reaction $J/\Psi \rightarrow \gamma p\bar{p}$. On the other hand, one has to concede that this enhancement as such does not provide any reliable additional support for the existence of the $X(1835)$ resonance, and likewise not for the existence of $p\bar{p}$ bound states or baryonia. With regard to the latter we want to remind the reader that our model calculations [12] as well as those of Loiseau and Wycech [16] are able to reproduce the $p\bar{p}$ spectrum. However, while the $N\bar{N}$ model used in [16] generates a near-threshold bound state (at $E \approx -5 - i50$ MeV) in the relevant 1S_0 state no such state is present in our 3S_0 amplitude (which describes the data equally well) and the one in the 1S_0 (at $E \approx -104 - i413$ MeV [23]) is too wide to have an influence in the physical region.

We believe that it will be rather difficult to establish experimentally a direct connection between the $X(1835)$ and the $p\bar{p}$ system. On the other hand, it would be still interesting to investigate whether this resonance is also visible in $p\bar{p}$ annihilation. The $X(1835)$ could be searched for in reactions like $p\bar{p} \rightarrow \pi^+ \pi^- X$, $X \rightarrow \pi^+ \pi^- \eta'$, etc. [24]. Measurements to get more information on these issues could be performed using the PANDA detector at




FIG. 2: Invariant FSI corrected $J/\Psi \rightarrow \gamma p\bar{p}$ amplitude $|A_{J/\Psi}|^2/|T|^2$ as a function of the $p\bar{p}$ mass. The squares show the values of $|A_{J/\Psi}|^2$ extracted from the BES data [3] via Eq. (2) and divided by the 1S_0 scattering amplitude squared for the $I = 0$ and $I = 1$ isospin states. The lines show the squared Breit-Wigner amplitudes for the $X(1835)$ [21], $\eta(1800)$ [22] and $\pi(1800)$ and $\eta(1760)$ [20] states.

the future FAIR project.

In this context let us also remind the reader that standard quark-model calculations like those in Ref. [25, 26, 27] do predict radial excitations of the η , η' around the $\eta(1800)$ mass. Thus, a very conventional interpretation of the structure found by the BES Collaboration should be also taken into consideration before speculating excessively on exotic explanations.

In summary, we have analyzed the near-threshold enhancement in the $p\bar{p}$ invariant mass spectrum from the $B \rightarrow K p\bar{p}$ decay reported recently by the BaBar Collaboration within the Jülich $N\bar{N}$ model. Our study shows that the mass dependence of the $p\bar{p}$ spectrum close to the threshold can be reproduced by the final state interaction. This explanation is in line with our previous

investigation of the $p\bar{p}$ invariant mass spectrum from the $J/\Psi \rightarrow \gamma p\bar{p}$ decay measured by the BES Collaboration.

Acknowledgments

We thank Mathew Graham for providing us with the most recent BaBar data. This work was partially supported by the DFG (SFB/TR 16, “Subnuclear Structure of Matter”), by the EU Integrated Infrastructure Initiative Hadron Physics Project (contract no. RII3-CT-2004-506078) and by Department of Energy under contract no. SURA-06-C0452.

-
- [1] K. Abe et al., Phys. Rev. Lett. **88**, 181803 (2002) [arXiv:hep-ex/0202017].
 - [2] K. Abe et al., Phys. Rev. Lett. **89**, 151802 (2002) [arXiv:hep-ex/0205083].
 - [3] J.Z. Bai et al., Phys. Rev. Lett. **91**, 022001 (2003) [arXiv:hep-ex/0303006].
 - [4] M.Z. Wang et al., Phys. Rev. Lett. **92**, 131801 (2004).
 - [5] S.B. Athar et al., Phys. Rev. D **73**, 032001 (2006).
 - [6] B. Aubert et al., Phys. Rev. D **72**, 051101 (R) (2005).
 - [7] see https://oraweb.slac.stanford.edu:8080/pls/slacquery/BABAR_DOCUMENTS.DetailedIndex?P_BP_ID=4092
 - [8] C.B. Dover and M. Goldhaber, Phys. Rev. D **15**, 1997 (1977).
 - [9] I.S. Shapiro, Phys. Rept. **35**, 129 (1978).
 - [10] C. K. Chua, W. S. Hou and S. Y. Tsai, Phys. Lett. B **544**, 139 (2002).
 - [11] J.L. Rosner, Phys. Rev. D **68**, 014004 (2003).
 - [12] A. Sibirtsev, J. Haidenbauer, S. Krewald, Ulf-G. Meißner and A.W. Thomas, Phys. Rev. D **71**, 054010 (2005) [arXiv:hep-ph/0411386].
 - [13] B. Kerbikov, A. Stavinsky, and V. Fedotov, Phys. Rev. C **69**, 055205 (2004) [arXiv:hep-ph/0402054].
 - [14] D.V. Bugg, Phys. Lett. B **598**, 8 (2004) [arXiv:hep-ph/0406293].
 - [15] B.S. Zou and H.C. Chiang, Phys. Rev. D **69**, 034004 (2004) [arXiv:hep-ph/0309273].
 - [16] B. Loiseau and S. Wycech, Phys. Rev. C **72**, 011001 (2005) [arXiv:hep-ph/0501112].
 - [17] T. Hippchen, J. Haidenbauer, K. Holinde, V. Mull, Phys. Rev. C **44**, 1323 (1991); V. Mull, J. Haidenbauer, T. Hippchen, K. Holinde, Phys. Rev. C **44**, 1337 (1991).
 - [18] V. Mull, K. Holinde, Phys. Rev. C **51**, 2360 (1995).
 - [19] E. Byckling and K. Kajantie, Particle Kinematics, John Wiley and Sons (1973).
 - [20] Particle Data Group, Phys. Lett. B **592**, 1 (2004).
 - [21] M. Ablikim et al., Phys. Rev. Lett. **95**, 262001 (2005) [arXiv:hep-ex/0508025].
 - [22] D. Bisello et al., Phys. Rev. D **39**, 701 (1989).
 - [23] Note that there is a misprint in the binding energy given in Ref. [12].
 - [24] E. Klempt, arXiv:hep-ex/0404270.
 - [25] D.P. Stanley and D. Robson, Phys. Rev. D **21**, 3180 (1980).
 - [26] S. Godfrey and N. Isgur, Phys. Rev. D **32**, 189 (1985).
 - [27] M. Koll, R. Ricken, D. Merten, B. C. Metsch and H. R. Petry, Eur. Phys. J. A **9**, 73 (2000) [arXiv:hep-ph/0008220].